

Mixing Instructions

With **KBPC 20 (code: 969553)**
 Country **Tanzania**
 Specie **Broiler Breeders**
 Batch weight **1,000 kg**



Version: 5/1/2020 Code	Ingredient	Unit (kg)	22,120 Starter 0-3 weeks	22,220 Grower 3-10 weeks	22,320 Developer 11-19 weeks	22,420 Pre-Breeder 19-24 weeks	22520 Breeder 1 24-45 weeks	22620 Breeder 2 >45 weeks	22900 Males >25 weeks
969553	KBPC 20		250	230	200	200	220	200	180
KN627C	MAIZE		570	550	520	585	620	600	530
KN699	SOYBEANMEAL 46CP/6CF		110	50					
KN673-1C	WHEAT BRAN		55	150	260	170	65	95	270
KN778C	LIMESTONES 30% CA		15	20	20	45	95	105	20
	Total		1000	1000	1000	1000	1000	1000	1000

CALCULATED ANALYSIS

001	Crude Protein	%	20.7	18.4	16.3	15.5	15.0	14.5	15.8
002	Crude Fat	%	4.2	4.2	4.2	4.1	3.9	3.9	4.2
003	Crude Fibre	%	4.6	5.0	5.6	4.7	3.8	3.9	5.6
089	Dry matter	%	87.8	87.8	87.8	88.0	88.6	88.7	87.8
004	Ash	%	5.7	6.0	5.8	7.9	12.5	13.4	5.7
012	ME Layer	Kcal	2870	2810	2740	2790	2750	2690	2750
011	ME Poultry	Kcal	2830	2760	2690	2740	2700	2650	2700
1750	Digestible Lysine	%	0.92	0.78	0.64	0.61	0.61	0.58	0.61
1768	Digestible Methion + Cystin	%	0.72	0.66	0.59	0.57	0.57	0.55	0.57
1753	Digestible Treonin	%	0.65	0.57	0.49	0.47	0.47	0.45	0.47
016	Total lysin	%	1.05	0.89	0.74	0.70	0.69	0.66	0.70
019	Total Methionin + Cystin	%	0.80	0.74	0.67	0.64	0.63	0.61	0.64
020	Total Treonin	%	0.79	0.69	0.60	0.58	0.57	0.55	0.58
048	Calcium	%	0.77	0.89	0.86	1.60	3.11	3.39	0.84
049	Total Phosphor	%	0.52	0.54	0.56	0.50	0.44	0.44	0.55
145	Digestible Phosphor	%	0.34	0.32	0.30	0.29	0.29	0.27	0.29
052	Sodium	%	0.21	0.19	0.17	0.17	0.18	0.17	0.15
050	Chloride	%	0.32	0.30	0.27	0.27	0.28	0.26	0.25
ZC	Coccidiostat		Not Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added
999	Essential vitamins & trace minerals		Added	Added	Added	Added	Added	Added	Added

Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- When oil is available; replace 0,5% corn for 0,5% oil. This can improve FCR and technical performance.
- Always make sure all birds have access to enough feed.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- Make sure birds always have access to fresh, cool and clean water.
- In case of management related issues, ask for the Koudijs Management Guide.

V10_02

Mixing Instructions											
With KBPC 20 (code: 969553)											
Country Tanzania											
Species Broiler Breeders											
Batch weight 1,000 kg											
Version: 5/1/2020											
Code Ingredient			Unit (kg)								
			0-3 weeks	3-10 weeks	11-19 weeks	19-24 weeks	24-45 weeks	>45 weeks	>25 weeks		
			21120 Starter	22220 Grower	22320 Developer	22420 Pre-Breeder	22520 Breeder 1	22620 Breeder 2	22900 Males		
969553 KBPC 20			250	230	200	200	220	200	180		
KN627C MAIZE			570	550	520	585	620	600	530		
KN699 SOYBEANMEAL 46CP6CF			110	90							
KN673-1C WHEAT BRAN			55	150	260	170	65	95	270		
KN776C LIMESTONES 30% CA			15	20	20	45	95	105	20		
Total			1000	1000	1000	1000	1000	1000	1000		

