

Mixing Instructions

With **969574 & 969452**
 Country **Tanzania**
 Specie **Broiler**
 Batch weight **1.000 kg**

KBC 35/30 Optimal



Version: 19-5-2023

Code Ingredient	Unit (kg)	Starter 0 - Starter 8-14 days	Grower 11 Grower 15-28 days	Finisher Finisher 29 days- till end
969574 Galdus	min 175gr/DOC			
KN628C MAIZE 7.5% CP		600	620	670
969452 KBC 35/30 SA		350	330	300
KN660C MAIZEBRAN		50	50	30
Total		1.000	1.000	1.000

CALCULATED ANALYSIS

001 Crude Protein	%	23,1	20,0	19,2	18,1
002 Crude Fat	%	5,3	4,2	4,2	4,1
003 Crude Fibre	%	4,0	3,2	3,1	3,0
089 Dry matter	%	89,4	87,3	87,3	87,2
004 Ash	%	6,0	5,6	5,4	5,0
098 ME Poultry	Kcal	3020	2980	2990	3040
1750 Digestible Lysine	%	1,27	1,13	1,08	1,00
1768 Digestible Methion + Cystin	%	0,86	0,76	0,74	0,70
048 Calcium	%	0,86	0,92	0,87	0,79
049 Total Phosphor	%	0,55	0,42	0,41	0,39
145 Digestible Phosphor	%	0,45	0,39	0,37	0,34
147 Available Phosphor	%	0,47	0,40	0,38	0,35
052 Sodium	%	0,20	0,16	0,16	0,14
Som 15 Phytase			Added	Added	Added
Som 16 NSP Enzyme			Not Added	Not Added	Not Added
ZC Coccidiostat			Added	Added	Added
999 Essential vitamins & trace minerals		Added	Added	Added	Added

Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
- For the best results; start with the Koudijs Galdus 2mm pre-starter (minimum 175 gram/bird total, preferably more).
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- In case of management related issues, ask for the Koudijs Management Guide.
- Make sure birds always have access to fresh, cool and clean water.
- For a better homogeneity of the feed in the vertical mixers, make sure that the mixing time is 3-4 minutes.

V10_08