

Mixing Instructions

With **969574 & 969452**
 Country **Tanzania**
 Specie **Broiler**
 Batch weight **1.000 kg**

KBC 35/30 Performance



Version: 19-5-2023	Galdus	301-9002 Starter	302-9002 Grower	303-9002 Finisher	
Code Ingredient	Unit (kg)	0-7 days	8-14 days	15-28 days	29 days- till end
969574 Galdus		min 175gr/DOC			
KN628C MAIZE 7.5% CP			645	655	680
969452 KBC 35/30 SA			350	330	300
KN768 SOYBEAN OIL			5	15	20
Total			1.000	1.000	1.000

CALCULATED ANALYSIS

001 Crude Protein	%	23,1	19,8	19,0	17,9
002 Crude Fat	%	5,3	4,4	5,4	5,9
003 Crude Fibre	%	4,0	3,0	2,9	2,9
089 Dry matter	%	89,4	87,3	87,4	87,4
004 Ash	%	6,0	5,5	5,3	4,9
098 ME Poultry	Kcal	3020	3050	3120	3180
1750 Digestible Lysine	%	1,27	1,13	1,07	0,99
1768 Digestible Methion + Cystin	%	0,86	0,76	0,73	0,69
048 Calcium	%	0,86	0,92	0,87	0,79
049 Total Phosphor	%	0,55	0,42	0,41	0,39
145 Digestible Phosphor	%	0,45	0,39	0,37	0,34
147 Available Phosphor	%	0,47	0,40	0,38	0,35
052 Sodium	%	0,20	0,16	0,16	0,14
Som 15 Phytase		Added	Added	Added	Added
Som 16 NSP Enzyme		Added	Not Added	Not Added	Not Added
ZC Coccidiostat		Not Added	Added	Added	Added
999 Essential vitamins & trace minerals		Added	Added	Added	Added

Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
- For the best results; start with the Koudijs Galdus 2mm pre-starter (minimum 175 gram/bird total, preferably more).
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- In case of management related issues, ask for the Koudijs Management Guide.
- Make sure birds always have access to fresh, cool and clean water.
- For a better homogeneity of the feed in the vertical mixers, make sure that the mixing time is 3-4 minutes.

V10_08