

## Mixing Instructions

With **969501**  
 Country **Tanzania**  
 Specie **Layers**  
 Batch weight **1.000 kg**

### Optimal Mixing (with oil) - Production KLC 5



Version: 14-11-2023		204-9208 Prelay	205-9208 Prepeak	206-9208 Layer 1	207-9208 Layer 2
Code Ingredient	Unit (kg)	19-20 weeks	21-28 weeks	29-55 weeks	56 weeks - end
KN628C MAIZE 7.5% CP		460	480	500	490
KN699 SOYBEANMEAL 46%CP/6%CF		130	140	165	160
969501 KLC 5 + CARO		50	50	50	50
KN660C MAIZEBRAN		110	70	40	40
KN673C WHEAT POLLARD/BRAN		110	70	35	35
KN718 SUNFLOWERCAKE		80	100	100	105
KN778-01 LIMESTONES 36 COURSE		55	80	95	105
KN770 VEGETABLE OIL		5	10	15	15
<b>Total</b>		<b>1.000</b>	<b>1.000</b>	<b>1.000</b>	<b>1.000</b>

### CALCULATED ANALYSIS

001 Crude Protein	%	15,8	15,8	16,3	16,1
002 Crude Fat	%	5,8	6,2	6,3	6,4
003 Crude Fibre	%	5,4	5,5	5,3	5,4
089 Dry matter	%	88,6	89,0	89,2	89,4
004 Ash	%	9,3	11,7	13,1	14,1
600k ME Poultry	Kcal	2710	2690	2690	2660
600 ME Poultry	MJ	11,34	11,24	11,25	11,12
1750 Digestible Lysine	%	0,68	0,69	0,73	0,72
1768 Digestible Methion + Cystin	%	0,64	0,64	0,65	0,64
048 Calcium	%	2,35	3,25	3,79	4,15
049 Total Phosphor	%	0,42	0,40	0,39	0,39
145 Digestible Phosphor	%	0,31	0,31	0,31	0,31
147 Available Phosphor	%	0,31	0,31	0,31	0,31
052 Sodium	%	0,15	0,15	0,15	0,16
Som 15 Phytase		Added	Added	Added	Added
Som 16 NSP Enzyme		Added	Added	Added	Added
ZC Coccidiostat		Not Added	Not Added	Not Added	Not Added
554 Mycotoxin binder		Not Added	Not Added	Not Added	Not Added
999 Essential vitamins & trace minerals		Added	Added	Added	Added

#### Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- In case of management related issues, ask for the Koudijs Management Guide.
- Make sure birds always have access to fresh, cool and clean water.
- Economy mixings are focused on a low price per kg of complete feed and an economical performance.

V10\_09