

Mixing Instructions

With **969574 & 969452**
 Country **Tanzania**
 Specie **Rearing & Layer**
 Batch weight **1.000 kg**

KLC 10 Economical



Version: 19-5-2023

Code	Ingredient	Unit (kg)	Galdus	201-9202 Starter	202-9202 Grower	203-9206 Developer	205-9206 Transition	206-9206 Phase 1	207-9206 Phase 2
			0-10 days	1-6 weeks	7-12 weeks	13-18 weeks	19-28 weeks	29-50 weeks	> 51 weeks
	969574 Galdus		min 100gr/DOC						
	969452 KBC 35/30 SA			350	300				
	969502 KLC 10					90	100	100	100
	KN628C MAIZE 7.5% CP			510	500	450	510	510	505
	KN660 MAIZE BRAN			140	200	200	70	50	50
	KN778 LIMESTONES 36 FINE					10			
	KN718 SUNFLOWERCAKE					100	110	110	110
	KN697 SOYBEANMEAL 44CP/7CF					50	80	85	85
	KN778-01 LIMESTONES 36 COURSE						80	95	100
	KN665C RICE POLISHINGS					100	50	50	50
	Total			1.000	1.000	1.000	1.000	1.000	1.000

CALCULATED ANALYSIS

001	Crude Protein	%	23,1	20,2	18,6	14,3	14,9	15,0	14,9
002	Crude Fat	%	5,3	3,9	4,0	5,7	5,2	5,1	5,1
003	Crude Fibre	%	4,0	4,4	4,9	9,1	7,1	6,9	6,9
089	Dry matter	%	89,4	87,3	87,3	88,2	89,0	89,2	89,3
004	Ash	%	6,0	5,5	4,9	5,3	11,9	13,4	13,9
098	ME Poultry	Kcal	3020	2830	2800	2680	2640	2610	2590
1750	Digestible Lysine	%	1,27	1,13	1,00	0,64	0,72	0,73	0,73
1768	Digestible Methion + Cystin	%	0,86	0,77	0,71	0,62	0,65	0,65	0,65
048	Calcium	%	0,86	0,92	0,79	0,79	3,24	3,78	3,96
049	Total Phosphor	%	0,55	0,42	0,38	0,48	0,44	0,44	0,44
145	Digestible Phosphor	%	0,45	0,39	0,34	0,30	0,32	0,32	0,32
147	Available Phosphor	%	0,47	0,40	0,35	0,30	0,32	0,32	0,32
052	Sodium	%	0,20	0,17	0,14	0,14	0,15	0,15	0,15
Som 15	Phytase		Added	Added	Added	Added	Added	Added	Added
Som 16	NSP Enzyme		Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added
ZC	Coccidiostat		Not Added	Added	Added	Not Added	Not Added	Not Added	Not Added
999	Essential vitamins & trace minerals		Added	Added	Added	Added	Added	Added	Added

Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
- For the best results; start with the Koudijs Galdus 2mm pre-starter (minimum 100 gram/bird total, preferably more).
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- In case of management related issues, ask for the Koudijs Management Guide.
- Make sure birds always have access to fresh, cool and clean water.
- For a better homogeneity of the feed in the vertical mixers, make sure that the mixing time is 3-4 minutes.

V10_08