

Mixing Instructions

With **969452 & 969502**
 Country **Tanzania**
 Specie **Rearing & Layer**
 Batch weight **1.000 kg**

KLC 10 Optimal



Version: 19-5-2023

Code	Ingredient	Unit (kg)	Galdus	201-9202	202-9202	203-9205	204-9205	205-9205	206-9205	207-9205
			0-10 days	Starter	Grower	Developer	Prelay	Prepeak	Phase 1	Phase 2
				1-6 weeks	7-12 weeks	13-17 weeks	18-19 weeks (till 5% production)	20-28 weeks	29-45 weeks	46 weeks - end
969574	Galdus	min 100gr/DOC								
969452	KBC 35/30 SA			350	300					
969502	KLC 10					90	100	100	100	100
KN628C	MAIZE 7.5% CP			510	500	500	550	560	565	555
KN660	MAIZE BRAN			140	200	250	135	80	50	50
KN697	SOYBEANMEAL 44CP/7CF					70	90	100	100	100
KN718	SUNFLOWERCAKE					70	70	80	90	90
KN778	LIMESTONES 36 FINE					20				
KN778-01	LIMESTONES 36 COURSE						55	80	95	105
Total				1.000	1.000	1.000	1.000	1.000	1.000	1.000

CALCULATED ANALYSIS

001	Crude Protein	%	23,1	20,2	18,6	14,5	15,0	15,2	15,1	15,1
002	Crude Fat	%	5,3	3,9	4,0	4,8	4,6	4,6	4,6	4,6
003	Crude Fibre	%	4,0	4,4	4,9	7,1	6,1	5,7	5,6	5,6
089	Dry matter	%	89,4	87,3	87,3	87,8	88,3	88,7	88,9	89,1
004	Ash	%	6,0	5,5	4,9	5,0	8,7	11,3	12,8	13,8
098	ME Poultry	Kcal	3020	2830	2800	2720	2720	2690	2680	2640
1750	Digestible Lysine	%	1,27	1,13	1,00	0,66	0,73	0,75	0,75	0,75
1768	Digestible Methion + Cystin	%	0,86	0,77	0,71	0,63	0,65	0,66	0,65	0,65
048	Calcium	%	0,86	0,92	0,79	0,96	2,24	3,15	3,69	4,05
049	Total Phosphor	%	0,55	0,42	0,38	0,35	0,36	0,37	0,38	0,38
145	Digestible Phosphor	%	0,45	0,39	0,34	0,28	0,31	0,31	0,31	0,31
147	Available Phosphor	%	0,47	0,40	0,35	0,28	0,31	0,31	0,32	0,32
052	Sodium	%	0,20	0,17	0,14	0,14	0,15	0,15	0,15	0,15
Som 15	Phytase	Added	Added	Added	Added	Added	Added	Added	Added	Added
Som 16	NSP Enzyme	Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added
ZC	Coccidiostat	Not Added	Added	Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added
999	Essential vitamins & trace minerals	Added	Added	Added	Added	Added	Added	Added	Added	Added

Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
- For the best results; start with the Koudijs Galdus 2mm pre-starter (minimum 100 gram/bird total, preferably more).
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- In case of management related issues, ask for the Koudijs Management Guide.
- Make sure birds always have access to fresh, cool and clean water.
- For a better homogeneity of the feed in the vertical mixers, make sure that the mixing time is 3-4 minutes.

V10_08