

Mixing Instructions

With **969574 & 969564**
 Country **Tanzania**
 Specie **Rearing & Layer**
 Batch weight **1.000 kg**

KLC 20 Economical



Version: 19-5-2023

Code	Ingredient	Unit (kg)	Galdus	201-9202 Starter	202-9202 Grower	203-9203 Developer	205-9203 Transition	206-9203 Phase 1	207-9203 Phase 2
			0-10 days	1-6 weeks	7-12 weeks	13-18 weeks	19-28 weeks	29-50 weeks	> 51 weeks
	969574 Galdus		min 100gr/DOC						
	969452 KBC 35/30 SA			350	300				
	969564 KLC 20 + Caro					180	200	200	200
	KN628C MAIZE 7.5% CP			510	500	450	500	520	510
	KN660 MAIZE BRAN			140	200	190	100	85	85
	KN718 SUNFLOWERCAKE					50	50	50	50
	KN778-01 LIMESTONES 36 COURSE						70	95	105
	KN665C RICE POLISHINGS					120	80	50	50
	KN778 LIMESTONES 36 FINE					10			
	Total			1.000	1.000	1.000	1.000	1.000	1.000

CALCULATED ANALYSIS

001	Crude Protein	%	23,1	20,2	18,6	15,2	15,2	15,0	14,9
002	Crude Fat	%	5,3	3,9	4,0	5,3	4,8	4,5	4,5
003	Crude Fibre	%	4,0	4,4	4,9	8,0	6,2	5,5	5,5
089	Dry matter	%	89,4	87,3	87,3	88,1	88,7	88,9	89,0
004	Ash	%	6,0	5,5	4,9	5,6	11,3	13,3	14,3
098	ME Poultry	Kcal	3020	2830	2800	2700	2660	2630	2600
1750	Digestible Lysine	%	1,27	1,13	1,00	0,66	0,69	0,68	0,68
1768	Digestible Methion + Cystin	%	0,86	0,77	0,71	0,61	0,62	0,62	0,61
048	Calcium	%	0,86	0,92	0,79	0,85	2,96	3,80	4,16
049	Total Phosphor	%	0,55	0,42	0,38	0,52	0,48	0,44	0,44
145	Digestible Phosphor	%	0,45	0,39	0,34	0,29	0,31	0,31	0,31
147	Available Phosphor	%	0,47	0,40	0,35	0,28	0,32	0,31	0,31
052	Sodium	%	0,20	0,17	0,14	0,13	0,15	0,15	0,15
Som 15	Phytase		Added	Added	Added	Added	Added	Added	Added
Som 16	NSP Enzyme		Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added
ZC	Coccidiostat		Not Added	Added	Added	Not Added	Not Added	Not Added	Not Added
999	Essential vitamins & trace minerals		Added	Added	Added	Added	Added	Added	Added

Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
- For the best results; start with the Koudijs Galdus 2mm pre-starter (minimum 100 gram/bird total, preferably more).
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- In case of management related issues, ask for the Koudijs Management Guide.
- Make sure birds always have access to fresh, cool and clean water.
- For a better homogeneity of the feed in the vertical mixers, make sure that the mixing time is 3-4 minutes.

V10_08