

Mixing Instructions

With 969452 & 969564
Country Tanzania
Specie Rearing & Layer
Batch weight 1.000 kg

KLC 20 Optimal



Version: 19-5-2023

Code	Ingredient	Unit (kg)	Galdus	201-9202	202-9202	203-9202	204-9202	205-9202	206-9202	207-9202
			0-10 days	Starter	Grower	Developer	Prelay	Prepeak	Phase 1	Phase 2
				1-6 weeks	7-12 weeks	13-17 weeks	18-19 weeks (till 5% production)	20-28 weeks	29-45 weeks	46 weeks - end
969574	Galdus	min 100gr/DOC								
969452	KBC 35/30 SA			350	300					
969564	KLC 20 + Caro					180	200	200	200	200
KN628C	MAIZE 7.5% CP			510	500	500	550	590	555	550
KN660	MAIZE BRAN			140	200	280	145	50	50	50
KN778	LIMESTONES 36 FINE					15				
KN718	SUNFLOWERCAKE					25	55	80	100	100
KN778-01	LIMESTONES 36 COURSE						50	80	95	100
	Total			1.000	1.000	1.000	1.000	1.000	1.000	1.000

CALCULATED ANALYSIS

001	Crude Protein	%	23,1	20,2	18,6	14,9	15,5	15,5	15,7	15,7
002	Crude Fat	%	5,3	3,9	4,0	4,3	4,5	4,7	4,9	4,8
003	Crude Fibre	%	4,0	4,4	4,9	5,9	5,4	4,9	5,3	5,3
089	Dry matter	%	89,4	87,3	87,3	87,5	88,1	88,6	88,9	89,0
004	Ash	%	6,0	5,5	4,9	4,5	8,3	11,4	12,9	13,4
098	ME Poultry	Kcal	3020	2830	2800	2740	2760	2760	2690	2680
1750	Digestible Lysine	%	1,27	1,13	1,00	0,64	0,69	0,70	0,70	0,70
1768	Digestible Methion + Cystin	%	0,86	0,77	0,71	0,61	0,64	0,64	0,64	0,64
048	Calcium	%	0,86	0,92	0,79	0,80	2,09	3,18	3,72	3,90
049	Total Phosphor	%	0,55	0,42	0,38	0,36	0,39	0,40	0,41	0,41
145	Digestible Phosphor	%	0,45	0,39	0,34	0,28	0,30	0,31	0,31	0,31
147	Available Phosphor	%	0,47	0,40	0,35	0,29	0,31	0,31	0,31	0,31
052	Sodium	%	0,20	0,17	0,14	0,14	0,15	0,15	0,15	0,15
Som 15	Phytase		Added	Added	Added	Added	Added	Added	Added	Added
Som 16	NSP Enzyme		Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added
ZC	Coccidiostat		Not Added	Added	Added	Not Added	Not Added	Not Added	Not Added	Not Added
999	Essential vitamins & trace minerals		Added	Added	Added	Added	Added	Added	Added	Added

Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
- For the best results; start with the Koudijs Galdus 2mm pre-starter (minimum 100 gram/bird total, preferably more).
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- In case of management related issues, ask for the Koudijs Management Guide.
- Make sure birds always have access to fresh, cool and clean water.
- For a better homogeneity of the feed in the vertical mixers, make sure that the mixing time is 3-4 minutes.

V10_08