

Mixing Instructions

With 969452 & 969564
Country Tanzania
Specie Rearing & Layers
Batch weight 1.000 kg

KLC 20 Performance



Version: 19-5-2023

Code	Ingredient	Unit (kg)	Galdus	201-9201 Starter	202-9201 Grower	203-9201 Developer	204-9201 Prelay	205-9201 Prepeak	206-9201 Phase 1	207-9201 Phase 2
			0-10 days	1-6 weeks	7-12 weeks	13-17 weeks	18-19 weeks (till 5% production)	20-28 weeks	29-45 weeks	46 weeks - end
	969574 Galdus	min. 100gr/DOC								
	969452 KBC 35/30 SA			350	300					
	969564 KLC 20 + Caro					180	200	200	200	200
	KN628C MAIZE 7.5% CP			520	500	500	530	530	525	515
	KN660 MAIZE BRAN			75	75	130	160	95	75	75
	KN673 WHEAT BRAN			50	120	150				
	KN768 SOYBEAN OIL			5	5	5	10	15	15	15
	KN778 LIMESTONES 36 FINE					15				
	KN718 SUNFLOWERCAKE					20	40	80	90	90
	KN778-01 LIMESTONES 36 COURSE						60	80	95	105
	Total			1.000	1.000	1.000	1.000	1.000	1.000	1.000

CALCULATED ANALYSIS

001	Crude Protein	%	23,1	20,3	19,1	15,3	15,2	15,5	15,5	15,4
002	Crude Fat	%	5,3	4,4	4,4	4,8	5,3	6,1	6,2	6,1
003	Crude Fibre	%	4,0	4,0	4,3	5,3	5,1	5,3	5,3	5,3
089	Dry matter	%	89,4	87,5	87,5	87,7	88,3	88,8	89,1	89,2
004	Ash	%	6,0	5,7	5,3	4,9	9,2	11,3	12,8	13,8
098	ME Poultry	Kcal	3020	2880	2820	2750	2770	2780	2750	2720
1750	Digestible Lysine	%	1,27	1,14	1,03	0,67	0,68	0,70	0,70	0,69
1768	Digestible Methion + Cystin	%	0,86	0,77	0,71	0,61	0,63	0,64	0,63	0,63
048	Calcium	%	0,86	0,92	0,80	0,81	2,45	3,18	3,72	4,08
049	Total Phosphor	%	0,55	0,45	0,47	0,45	0,38	0,40	0,40	0,40
145	Digestible Phosphor	%	0,45	0,40	0,36	0,29	0,30	0,30	0,30	0,30
147	Available Phosphor	%	0,47	0,41	0,36	0,28	0,31	0,31	0,31	0,31
052	Sodium	%	0,20	0,16	0,14	0,13	0,15	0,15	0,15	0,15
Som 15	Phytase	Added	Added	Added	Added	Added	Added	Added	Added	Added
Som 16	NSP Enzyme	Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added
ZC	Coccidiostat	Not Added	Added	Added	Added	Not Added	Not Added	Not Added	Not Added	Not Added
999	Essential vitamins & trace minerals	Added	Added	Added	Added	Added	Added	Added	Added	Added

Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
- For the best results; start with the Koudijs Galdus 2mm pre-starter (minimum 100 gram/bird total, preferably more).
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- In case of management related issues, ask for the Koudijs Management Guide.
- Make sure birds always have access to fresh, cool and clean water.
- For a better homogeneity of the feed in the vertical mixers, make sure that the mixing time is 3-4 minutes.

V10_08